TABLE 4.7.3 SOUTHEAST REGION SEA TURTLE BYCATCH BY FISHERY (2013) Bycatch estimates are for live and dead releases in all fisheries. Estimates are included only for species with documented bycatch in a given year or years.¹

COMMON NAME	SCIENTIFIC NAME	YEAR	BYCATCH	UNIT	CV	FOOTNOTE(S)				
Atlantic and Gulf of Mexico HMS Pelagic Longline										
Leatherback sea turtle	Dermochelys coriacea	2013	365.60	INDIVIDUAL	.16					
Loggerhead sea turtle	Caretta caretta	2013	377.10	INDIVIDUAL	.16					
Fishery Total			742.70							
Gulf of Mexico Reef Fish Bottom Longline										
Loggerhead sea turtle	Caretta caretta	2013	11.90	INDIVIDUAL	.7					
Fishery Total			11.90							
Grand Total			754.60							

SOUTHEAST REGION SEA TURTLE BYCATCH BY STOCKS AND SPECIES

COMMON NAME	SCIENTIFIC NAME	YEAR	ВҮСАТСН	UNIT	FOOTNOTE(S)
Leatherback sea turtle	Dermochelys coriacea	2013	365.60	INDIVIDUAL	
Loggerhead sea turtle	Caretta caretta	2013	389.00	INDIVIDUAL	
GRAND TOTAL			754.60	INDIVIDUAL	

FOOTNOTE:

1 New estimates were not calculated for the shrimp trawl fisheries this year, but all four species (leatherback, loggerhead, green, and Kemp's ridley) are known to interact with shrimp trawls. NMFS published a Biological Opinion dated April 18, 2014, which contains more current sea turtle estimates for those fisheries.