Stock Name: Scalloped hammerhead - Atlantic Assessed in: October 2009

Parameter Name	Catch
Туре	Total Landings
Source	Observed
Basis	Numbers
Range	All
Statistic	Actual
Scale	1
Year	
1981	7367
1982	49626
1983	22814
1984	8490
1985	45529
1986	6808
1987	7600
1988	6194
1989	7580
1990	19536
1991	10160
1992	9649
1993	22330
1994	4554
1995	5741
1996	2652
1997	5884
1998	1959
1999	1275
2000	6695
2001	2602
2002	7724
2003	5896
2004	2585
2005	4135

TIME SERIES HEADER DESCRIPTIONS

Type: Provides a more detailed definition of the data being entered.

Source: Describes where a particular type of data comes from. Typical data sources include Model (output from an assessment model), Survey (index of survey observations), or Fishery (e.g. reported catch rather than a model estimate of catch).

Basis: Describes the units for the values being reported. For example: biomass-mt means stock weight in metric tons.

Range: Used in conjunction with type to refine the description of the data being entered. The range specifies a subset of the population to which the data apply. For example, Age 3+ means fish that are age 3 and older, or mature means just the mature portion of the stock.

Statistic: Describes the statistical characteristics of a time series column, and may include mean, median, index, observed, official, MCMC, lower 95% CI, upper 95% CI, etc.

Scalar: Describes a multiplier by which the reported values should be multiplied to restore them to their natural units. For example, if biomass is reported in 1000 mt, then a value of 1000 is entered in this field.