NOAA NMFS Stock Assessment Time Series Data

Stock Name: American Samoa Bottomfish Multi-species Complex

Assessed in: March 2015

Parameter Name	Catch	Abundance
Type	Biomass - Ibs	Biomass - Ibs
Source	Model	Model
Basis	Catch - Commercial and Recreational	Biomass - Exploitable
Range	All	All
Statistic	Point Estimate	Mean
Scale	1	1000
Year		
1986	64587	540
1987	19628	548
1988	33726	659
1989	32647	609
1990	11332	559
1991	1310	521
1992	9985	513
1993	14554	555
1994	33845	588
1995	27699	639
1996	30808	707
1997	32308	641
1998	12413	626
1999	15857	631
2000	19816	662
2001	37847	658
2002	34149	574
2003	19199	635
2004	17206	613
2005	16329	555
2006	7913	517
2007	21874	516
2008	34812	532
2009	47458	556
2010	9509	547
2011	26277	624
2012	13110	683
2013	23630	661

TIME SERIES HEADER DESCRIPTIONS

NOAA NMFS Stock Assessment Time Series Data

Type: Provides a more detailed definition of the data being entered.

Source: Describes where a particular type of data comes from. Typical data sources include Model (output from an assessment model), Survey (index of survey observations), or Fishery (e.g. reported catch rather than a model estimate of catch).

Basis: Describes the units for the values being reported. For example: biomass-mt means stock weight in metric tons.

Range: Used in conjunction with type to refine the description of the data being entered. The range specifies a subset of the population to which the data apply. For example, Age 3+ means fish that are age 3 and older, or mature means just the mature portion of the stock.

Statistic: Describes the statistical characteristics of a time series column, and may include mean, median, index, observed, official, MCMC, lower 95% CI, upper 95% CI, etc.

Scalar: Describes a multiplier by which the reported values should be multiplied to restore them to their natural units. For example, if biomass is reported in 1000 mt, then a value of 1000 is entered in this field.