Stock Name: Yellowtail flounder - Georges Bank

Assessed in: July 2015

Parameter Name	Index	Index	Catch
Туре	NEFSC Spring Survey	NEFSC Fall Survey	Total catch
Source	Survey	Survey	Fishery
Basis	Biomass-kg/tow	Biomass-kg/tow	Biomass-mt
Range	All	All	All
Statistic	Mean	Mean	Total
Scale	1	1	1
Year			
1963		12.8	16690
1964		13.6	19814
1965		9.1	19448
1966		3.9	13741
1967		7.7	15307
1968	2.8	10.5	18321
1969	11.2	9.8	21271
1970	5.1	5.0	21410
1971	4.6	6.4	15610
1972	6.5	6.3	18039
1973	2.9	6.5	16953
1974	2.7	3.7	17211
1975	1.7	2.3	16750
1976	2.3	1.5	14988
1977	1.0	2.8	10639
1978	.7	2.3	6944
1979	1.3	1.5	6935
1980	4.5	6.6	7539
1981	2.0	2.6	6979
1982	2.5	2.3	12520
1983	2.6	2.1	11989
1984	1.6	.6	6280
1985	1.0	.7	3267
1986	.8	.8	3474
1987	.3	.5	3580
1988	.6	.2	2759
1989	.7	1.0	1783
1990	.7	.7	4089
1991	.6	.7	2564
1992	1.6	.6	5299
1993	.5	.5	4300

NOAA NMFS Stock Assessment Time Series Data

1994	.7	.9	4158
1995	2.6	.4	1135
1996	2.9	1.3	1700
1997	4.4	3.8	2464
1998	2.3	4.3	3985
1999	9.3	8.0	4963
2000	6.7	5.8	7341
2001	5.0	11.6	7419
2002	9.6	3.8	5663
2003	6.7	4.0	6562
2004	1.9	5.1	6815
2005	3.4	2.5	3852
2006	2.4	4.5	2057
2007	4.7	8.2	1664
2008	3.2	7.1	1499
2009	4.9	6.7	1806
2010	5.9	2.2	1170
2011	2.6	2.5	1171
2012	4.0	2.5	725
2013	1.1	.9	218
2014	.7	1.0	159
2015	.5		

TIME SERIES HEADER DESCRIPTIONS

Type: Provides a more detailed definition of the data being entered.

Source: Describes where a particular type of data comes from. Typical data sources include Model (output from an assessment model), Survey (index of survey observations), or Fishery (e.g. reported catch rather than a model estimate of catch).

Basis: Describes the units for the values being reported. For example: biomass-mt means stock weight in metric tons.

Range: Used in conjunction with type to refine the description of the data being entered. The range specifies a subset of the population to which the data apply. For example, Age 3+ means fish that are age 3 and older, or mature means just the mature portion of the stock.

Statistic: Describes the statistical characteristics of a time series column, and may include mean, median, index, observed, official, MCMC, lower 95% CI, upper 95% CI, etc.

NOAA NMFS Stock Assessment Time Series Data

Scalar: Describes a multiplier by which the reported values should be multiplied to restore them to their natural units. For example, if biomass is reported in 1000 mt, then a value of 1000 is entered in this field.