

National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

JPL Activities

Task 1: Sounder Science Status


Task 2.3: Boundary Layer Science

**Eric Fetzer, Joao Teixeira, Thomas Pagano
and Kevin Bowman**

**Jet Propulsion Laboratory/California Institute of
Technology**

CLARREO Workshop
Newport News, Virginia
May 1, 2008

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Some Quotes

- “It’s easy to find problems with climate models.”
 - Anonymous Modeler
- “Models summarize our understanding of physical processes.”
 - Phil Rasch




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Some soapbox thinking...

- *Process scales* are important. Why?
 1. Climate change over the coming decades will be at process scales.
 2. Climate models represent processes (see previous quotes).
- The largest uncertainties in climate prediction are cloud feedbacks (especially low clouds).
 - Cloud scales are much smaller than 100 km.
 - I'll use this as an example; similar arguments may hold for other feedbacks.




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Climate Science Using Sounder Observations

- Measurements proposed for CLARREO in Decadal Survey are similar to those of IR sounders.
- What questions are the sounders addressing today?
- What key questions remain for the hyperspectral infrared?
- The future.

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

What issues are AIRS & TES addressing today? (publication 1st authors)

- Climate processes and model validation
 - Water vapor, temperature:
 - **Hurricanes, deep convection** (Atlas, Wu, Wong, Holloway)
 - **Lower & upper trop hydration** (Worden, Dessler, Ray)
 - **Supersturation** (Gettelman)
 - **IPPC models** (Pierce)
 - **Polar water cycle** (Ye, Gettelman)
 - **Climate feedbacks** (Su, Dessler)
 - Clouds
 - **Trends** (Sund)
 - **Microphysics** (Kahn, Nasiri)
 - **Inter-instrument consistency** (Kahn)

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

More issues AIRS & TES are addressing today

- Minor gases
 - **Ozone** (Bowman, Barnet)
 - **CO** (Osterman, McMillan)
 - **Methane** (Barnet)
 - **CO₂** (McNally, Morel, Chahine, Strow)
- Air-surface interaction
 - **Boundary layer over ocean** (Fetzer)
 - **Boundary layer over land** (Santanello)
- Radiative Balance
 - **IPCC models** (Huang)
 - **Water vapor radiative cooling** (Feldman)
 - **Cloud and aerosol longwave forcing** (Huang, de Souza)

Eric.J.Fetzer@jpl.nasa.gov



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Hyperspectral Sounder Retrievals and Spectra for Climate Model Validation

- Method 1: Data Products (L3)
 - The models are drier than AIRS observations by 10%-25% in the tropics below 800 hPa.
 - The models are more moist by 25%-100% between 300 and 600 hPa, especially in the extra-tropics.


** David W. Pierce, Tim P. Barnett, Eric J. Fetzer, Peter J. Gleckler, Three-dimensional tropospheric water vapor in coupled climate models compared with observations from the AIRS satellite system, GRL, VOL. 33, L21701, doi:10.1029/2006GL027060, 2006*


- Method 2. Radiances (L1b)
 - OLR Agrees with Models
 - Compensating Errors
 - Models dry in lower troposphere compensated by higher surface flux

** Huang et al. 2007.*



Unit: W m ⁻²	OLR		Window band	
	Total sky	Clear sky	Total sky	Clear sky
CERES	241.73	275.87	66.94	83.28
AM2	240.63	263.43	73.99	87.56
AM2-CERES	-1.10	-12.44	7.05	4.28






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


IPCC model mean water vapor is biased

Similarly temperature (not shown).

Mean Climatologies of AIRS and 17 IPCC Models.
From Pierce et al. 2006, GRL



Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

MODIS cannot resolve High Ice Clouds over Low Water Clouds

Standard Simulations



Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

AIRS IR has cloud info not available with MODIS.

Standard Simulations



Potentially Mixed-Phase Simulations




Water cloud curves are solid, ice cloud curves are dashed or dotted.

The biggest issue is the (large) 15 km scale of the AIRS footprint.
“Borrowed” from Shaima Nasiri.

~0.5 K phase separation


Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

AIRS data are mature enough to start deducing trends



Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


SE Pacific

Weaker annual cycle with height.




Temperature

Strong annual cycle at all heights.



Water vapor

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


AIRS CO₂ work continues to make progress

AIRS CO₂ Product (Chahine)


AIRS Mid-Tropospheric CO₂. July 2003, V5 Day 16 x 31




CO₂ Trending using Spectra (Strow)



AIRS Zonal CO₂ Trends (Strow)



Demonstrates <10mK/year Stability³




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Summary: Hyperspectral Sounders

- Sounders observe key climate signatures, especially at scales >50 km.
- Many of these processes are poorly represented in models (or not analyzed, yet).
- Trending is only beginning.
 - Climate prediction will improve with record length.
- What additional questions will CLARREO answer?
 - especially given IPCC 2007 concerns with cloud feedbacks.

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Boundary Layer Science

- Why?
 - IPCC: “Cloud feedbacks remain the largest source of uncertainty.”
 - Large uncertainties in the hydrologic cycle arise from poorly known land-surface interactions.





National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Low cloud climate feedbacks can be of opposite sign in IPCC models!

Models ranked by double CO_2 response.



Fewer low clouds with warming




More low clouds with warming

B.Soden/G.Stephens

GFDL and NCAR have opposite low cloud cover sensitivity to CO_2 doubling

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Remember This?

Spatial or Temporal Averaging Will Reduce Sensitivity to Clouds



King-Fai, Duane Waliser, Yuk Yung, CalTech

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

What horizontal scales control stratocumulus clouds in the PBL?


GOES-10 VISIBLE IMAGE AND DETAIL
October 17 2001, 15:00 UTC



Bretherton et al (2004)

Eric.J.Fetzer@jpl.nasa.gov

Scales from 1 to 100 km probably play a significant role





National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Climate models will have very high resolution in 10 years

- Global Cloud Resolving Models (1-10 km) exist and produce realistic simulations of clouds.
- Chapter 9 of Decadal Survey calls for high spatial resolution climate models, especially regionally.
- CLARREO resolution should be commensurate.




Miura et al. 2007

NICAM global model (3.5 km horizontal resolution):

(A) IR image; (B) model simulation

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Hyperspectral sounders have sufficient resolution for PBL T & q.

*But... They do not observe at
the scale of cloud processes.*




Frequency of temperature inversions
1-15 January 2003

1000-700 hPa water vapor (mm)

700-500 hPa water vapor (mm)

Occurrence Frequency of Temperature Inversions
and Water Vapor Distribution from AIRS. Figure 3
from Fetzer et al. 2004

Eric.J.Fetzer@jpl.nasa.gov



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Some Testable Hypotheses from increasing CO₂

- A shift in subtropical low clouds will occur over the next century, manifested at scales of 1-50 km.
 - The signal will be robust in the IR because of local changes in cloud microphysics.
 - The signal will be robust in the visible because of changes in local reflectivity.
 - The energetics will be much stronger than local CO₂ forcing.
 - Associate clouds feedbacks may dominate the climate change signal.
- Increased Walker Cell strength will lead to changes in the tropical Pacific subvisible cirrus.
 - The signal will detectable in the IR through cloud microphysics.
 - The energetics will be like local CO₂ forcing.

**We think CLARREO should address the key issue in climate:
Cloud Feedbacks**

Eric.J.Fetzer@jpl.nasa.gov




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Low Cloud Summary

- 1) Low cloud climate feedbacks are presently a key problem in climate change prediction
- 2) Stratocumulus clouds are the major players in cloud radiative forcing and are behind low-cloud climate feedbacks
- 3) Mesoscale physical processes (between 1 to 100 km) play a significant role in controlling existence of stratocumulus
- 4) High-resolution (1-10 km) models are becoming more frequent and are “intriguingly” realistic in stratocumulus simulations
- 5) Observations at these resolutions are necessary to understand low cloud climate feedbacks and to validate future models



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

What We Will Do

- Examine spatial variability of radiances from TES, AIRS and MODIS associated with:
 - Stratocumulus
 - Water vapor
 - Cirrus
- Examine the same in high resolution models.
 - Start with WRF.
- What are the dominant scales CLARREO should observe?
 - to best exploit simultaneous, multispectral information.

Eric.J.Fetzer@jpl.nasa.gov